

# Probability

- If  $E$  and  $F$  are two events associated with the sample space of a random experiment, then the conditional probability of event  $E$ , given that  $F$  has already occurred, is denoted by  $P(E/F)$  and is given by the formula:

$$P(E/F) = \frac{P(E \cap F)}{P(F)}, \text{ where } P(F) \neq 0$$

## Example:

A die is rolled twice and the sum of the numbers appearing is observed to be 7. What is the conditional probability that the number 3 has appeared at-least once?

## Solution:

Let  $E$ : Event of getting the sum as 7 and  $F$ : Event of appearing 3 at-least once

Then  $E = \{(1, 6), (2, 5), (3, 4), (4, 3), (5, 2), (6, 1)\}$  and

$$F = \{(1, 3), (2, 3), (3, 3), (4, 3), (5, 3), (6, 3), (3, 1), (3, 2), (3, 4), (3, 5), (3, 6)\}$$

$$\therefore E \cap F = \{(3, 4), (4, 3)\}$$

$$n(E) = 6, n(F) = 11 \text{ and } n(E \cap F) = 2$$

$$P\left(\frac{F}{E}\right) = \frac{P(E \cap F)}{P(E)} = \frac{n(E \cap F)}{n(E)} = \frac{2}{6} = \frac{1}{3}$$

- If  $E$  and  $F$  are two events of a sample space  $S$  of an experiment, then the following are the properties of conditional probability:
  - $0 \leq P(E/F) \leq 1$

- $P(F/F) = 1$
- $P(S/F) = 1$
- $P(E'/F) = 1 - P(E/F)$
- If  $A$  and  $B$  are two events of a sample space  $S$  and  $F$  is an event of  $S$  such that  $P(F) \neq 0$ , then
  - $P((A \cup B)/F) = P(A/F) + P(B/F) - P((A \cap B)/F)$
  - $P((A \cup B)/F) = P(A/F) + P(B/F)$ , if the events  $A$  and  $B$  are disjoint.

• **Multiplication theorem of probability:** If  $E$ ,  $F$ , and  $G$  are events of a sample space  $S$  of an experiment, then

- $P(E \cap F) = P(E) \cdot P(F/E)$ , if  $P(E) \neq 0$
- $P(E \cap F) = P(F) \cdot P(E/F)$ , if  $P(F) \neq 0$
- $P(E \cap F \cap G) = P(E) \cdot P(F/E) \cdot P(G/(E \cap F)) = P(E) \cdot P(F/E) \cdot P(G/EF)$

- Two events  $E$  and  $F$  are said to be independent events, if the probability of occurrence of one of them is not affected by the occurrence of the other.
- If  $E$  and  $F$  are two independent events, then
  - $P(F/E) = P(F)$ , provided  $P(E) \neq 0$
  - $P(E/F) = P(E)$ , provided  $P(F) \neq 0$
- If three events  $A$ ,  $B$ , and  $C$  are independent events, then
 
$$P(A \cap B \cap C) = P(A) \cdot P(B) \cdot P(C)$$
- If the events  $E$  and  $F$  are independent events, then
  - $E'$  and  $F$  are independent
  - $E'$  and  $F'$  are independent
- A set of events  $E_1, E_2, \dots, E_n$  is said to represent a partition of the sample space  $S$ , if
  - $E_i \cap E_j = \emptyset, i \neq j, i, j = 1, 2, 3, \dots, n$

- $E_1 \cup E_2 \cup \dots \cup E_n = S$
- $P(E_i) > 0, \forall i = 1, 2, 3, \dots, n$

- **Bayes' Theorem:** If  $E_1, E_2, \dots, E_n$  are  $n$  non-empty events, which constitute a partition of sample space  $S$ , then

$$P(E_i / A) = \frac{P(E_i)P(A/E_i)}{\sum_{j=1}^n P(E_j)P(A/E_j)}, i = 1, 2, 3, \dots, n$$

**Example:**

There are three urns. First urn contains 3 white and 2 red balls, second urn contains 2 white and 3 red balls, and third urn contains 4 white and 1 red balls. A white ball is drawn at random. Find the probability that the white ball is drawn from the third urn?

**Solution:**

Let  $E_1, E_2$  and  $E_3$  be the events of choosing the first second and third urn respectively.

Then,  $P(E_1) = P(E_2) = P(E_3) = \frac{1}{3}$

Let  $A$  be the event that a white ball is drawn.

Then,  $P\left(\frac{A}{E_1}\right) = \frac{3}{5}, P\left(\frac{A}{E_2}\right) = \frac{2}{5}$  and  $P\left(\frac{A}{E_3}\right) = \frac{4}{5}$

By the theorem of total probability,

$$\begin{aligned}
P(A) &= P(E_1) \times P\left(\frac{A}{E_1}\right) + P(E_2) \times P\left(\frac{A}{E_2}\right) + P(E_3) \times P\left(\frac{A}{E_3}\right) \\
&= \frac{1}{3} \times \frac{3}{5} + \frac{1}{3} \times \frac{2}{5} + \frac{1}{3} \times \frac{4}{5} \\
&= \frac{3}{5}
\end{aligned}$$

By Bayes' theorem,

probability of getting the ball from third urn given that it is white

$$= P\left(\frac{E_3}{A}\right) = \frac{P(E_3)P\left(\frac{A}{E_3}\right)}{P(A)} = \frac{\frac{1}{3} \times \frac{4}{5}}{\frac{3}{5}} = \frac{4}{9}$$

- A random variable is a real-valued function whose domain is the sample space of a random experiment.
- The probability distribution of a random variable  $X$  is the system of numbers:

|         |       |       |         |       |
|---------|-------|-------|---------|-------|
| $X:$    | $x_1$ | $x_2$ | $\dots$ | $x_n$ |
| $P(X):$ | $p_1$ | $p_2$ | $\dots$ | $p_n$ |

Where,  $P_i > 0 = \sum_{i=1}^n p_i = 1, i = 1, 2, \dots, n$

Here, the real numbers  $x_1, x_2, \dots, x_n$  are the possible values of the random variable  $X$  and  $p_i$  ( $i = 1, 2, \dots, n$ ) is the probability of the random variable  $X$  taking the value of  $x_i$  i.e.,  $P(X = x_i) = p_i$

- **Mean/expectation of a random variable:** Let  $X$  be a random variable whose possible values  $x_1, x_2, x_3 \dots x_n$  occur with probabilities  $p_1, p_2, p_3 \dots p_n$  respectively. The mean of  $X$  (denoted by  $m$ ) or the expectation of

$X$  (denoted by  $E(X)$ ) is the number  $\sum_{i=1}^n x_i p_i$ .

That is,  $E(X) = \mu = \sum_{i=1}^n x_i p_i = x_1 p_1 + x_2 p_2 + \dots + x_n p_n$

- **Variance of a random variable:** Let  $X$  be a random variable whose possible values  $x_1, x_2 \dots x_n$  occur with probabilities  $p(x_1), p(x_2) \dots p(x_n)$  respectively. Let  $m = E(X)$  be the mean of  $X$ . The variance of  $X$  denoted by  $\text{Var}(X)$  or  $\sigma_x^2$  is calculated by any of the following formulae:

- $\sigma_x^2 = \sum_{i=1}^n (x_i - \mu)^2 p(x_i)$

- $\sigma_x^2 = E(X - \mu)^2$

- $\sigma_x^2 = \sum_{i=1}^n x_i^2 p(x_i) - \left[ \sum_{i=1}^n x_i p(x_i) \right]^2$

- $\sigma_x^2 = E(X^2) - [E(X)]^2$  where  $[E(X)]^2 = \sum_{i=1}^n x_i^2 p(x_i)$

It is advisable to students to use the fourth formula.

- **Binomial distribution:** For binomial distribution  $B(n, p)$ , the probability of  $x$  successes is denoted by  $P(X = x)$  or  $P(X)$  and is given by  $P(X = x) = {}^n C_x q^{n-x} p^x$ ,  $x = 0, 1, 2, \dots, n$ ,  $q = 1 - p$   
Here,  $P(X)$  is called the probability function of the binomial distribution.

### Example:

An unbiased coin is tossed 5 times. Find the probability of getting at least 4 heads.

### Solution:

Let the random variable  $X$  denotes the number of heads.

Here,  $n = 5$  and  $P$  (getting a head)  $= \frac{1}{2}$

$$\therefore p = \frac{1}{2} \text{ and } q = 1 - \frac{1}{2} = \frac{1}{2}$$

$$P(X = r) = {}^n C_r p^r q^{n-r} = {}^5 C_r \left(\frac{1}{2}\right)^r \left(\frac{1}{2}\right)^{5-r} = {}^5 C_r \left(\frac{1}{2}\right)^5$$

$P$  (getting at-least 4 heads)

$$= P(X \geq 4)$$

$$= P(X = 4) + P(X = 5)$$

$$= {}^5C_4 \left(\frac{1}{2}\right)^5 + {}^5C_4 \left(\frac{1}{2}\right)^5$$

$$= (5 + 1) \left(\frac{1}{2}\right)^5$$

$$= 6 \times \frac{1}{32}$$

$$= \frac{3}{16}$$